

Isolation and Molecular Characterization of Dermatophytes in Donkeys

Elham Abdelbasit Suleiman^{1*}, Wisal Gaafar Abdalla¹, Ahmed Haroon Ahmed²
and Mohamed Awad Abdelgabar¹

¹Central Veterinary Research Laboratory, Department of Mycology, Animal Resources Research Corporation, P.O. Box 8067 (El Amarat), Khartoum, Sudan.

²Eldamazine Regional Veterinary Research Laboratory, Sudan.

Authors' contributions

This work was carried out in collaboration between all authors. Author EAS designed the study, wrote the protocol and the first draft of the manuscript. Author WGA and EAS managed the analyses of the study. Author AHA managed the literature searches. All authors shared the laboratory work read and approved the final manuscript.

Article Information

DOI: 10.9734/IJTDH/2017/36425

Editor(s):

(1) Ravindra Nath Sharma, Department of Pathobiology, School of Veterinary Medicine, St. George's University, Grenada, West Indies.

Reviewers:

(1) Eliton da Silva Vasconcelos, Federal University of São Carlos, Brazil.

(2) Daisy Machado, Universidade Estadual de Campinas, Brazil.

Complete Peer review History: <http://www.sciencedomain.org/review-history/21370>

Original Research Article

Received 26th August 2017
Accepted 20th September 2017
Published 12th October 2017

ABSTRACT

Background: Dermatophytosis is a fungal infection of skin, hair and nail caused by dermatophytes. The zoonotic nature of the disease requires early detection for implication of treatment.

Aim: The study was conducted to investigate into the cause of skin infection among donkeys in Eldamazine State, Central Sudan.

Study Design: This a case history studies over one year duration.

Methodology: A total of 31 samples of skin scrapings from donkeys with an age ranged 2-20years, from Eldamazine state, Sudan, were collected. Samples were cultured onto Sabouraud's (SDA) media. The developed colonies were identified by convention method and characterized by molecular techniques using polymerase chain reaction.

Results: *Trichophyton mentagrophytes* and *Trichophyton verrucosum* were the predominant isolated dermatophytes revealed on investigation. Amplification of the β -tubulin gene, by polymerase chain reaction, and analysis of the amplicon sequence, further confirmed their

*Corresponding author: Email: elham842001@yahoo.com;

identification. The study revealed first report on characterization of these isolates by molecular methods using β - tubulin primers.

Conclusion: The present investigation showed that the integration of different methods and techniques led to identification of useful molecular marker for standardization of taxonomical studies of *dermatophyte* species.

Keywords: *Dermatophytes; beta tubulin primers; conventional methods; PCR.*

1. INTRODUCTION

Dermatophytosis (ringworm or tinea), is a superficial skin infection caused by closely related keratinophilic fungi [1]. They have the ability to degrade keratin and invade the skin and its appendages [2]. Ringworm is a major Public and Animal Health problem in various regions of the world resulting in great economic loss [3]. *T. mentagrophytes* var. *erinacei* accommodates the anthropophilic and zoophilic strains [4]. In man, *T. mentagrophytes* and *Microsporum canis* were reported to be the most common dermatophytes responsible for tinea infections in man [5,6]. Animals can get infection from soil while digging, rolling, and lying down. Infection depends on the fungal species, age, and health, condition of exposed skin surfaces, grooming behavior, and nutrition of the animal. However, broken hairs with associated spores are important sources for spread of the disease [7]. The identification of dermatophytes is based on methods that focus on morphological, physiological, ecological, and genetic characteristics [8,9,10].

1.1 Prevalence of Dermatophytosis in Donkeys

Donkeys (*Equus asinus*) are of great economic value being used for income generating activities. They can be used for riding, pack transport, pulling carts and for farm work [11]. Pack donkeys are an actual promising choice for transport in third world countries [12]. Despite their great value, they are unfortunately not given enough veterinary attention although they suffer many health problems and diseases such as dermatophytosis affecting their viability and ability to work. Dermatophytes are known to grow best in warm and humid environments and are therefore more common in tropical and subtropical regions and may present in epidemic proportions in areas with high rates of humidity but the geographic distribution varies according to the species [13,14]. Reports on ringworm in donkeys are scanty and refer to zoonotic agents such as *T. mentagrophytes* and

T. verrucosum. Recently *Microsporum racemosum* has been isolated from naturally infected donkeys [15]. In the Sudan, *T. verrucosum* has been isolated from donkeys ringworm [16]. Additionally, a severe outbreak of ringworm among 69 adult domestic donkeys in Sudan was described due to *T. mentagrophytes* [17]. However, in horses infections are common. *Trichophyton equinum* and *T. mentagrophytes* were identified as the primary cause of ringworm in horses [18,10]. *M. gypseum*, *M. canis*, and *T. verrucosum* have also been isolated [19,6]. Moreover, *Trichophyton* and *Microsporum* species associated with apparently normal horses' hair from Riyadh, KSA were isolated [20]. Recently, *T. mentagrophytes* var. *mentagrophytes* causing Ringworm in Horses in Al Ahsa Province, Kingdom of Saudi Arabia has been isolated [21].

2. MATERIALS AND METHODS

2.1 Clinical Examination

Thirty-one donkeys from Eldamazine city, central of the Sudan, in the age of 2-20 years, clinically suspected as having dermatophytosis were randomly selected for this study.

Skin lesions of suspected animals were examined and a complete clinical examination of all affected animals, was performed. The shape, size, position and distribution of lesions were observed.

2.2 Mycological Investigations

2.2.1 Macro and microscopic study

These included collection of samples, direct examination and culture. Affected areas were cleaned and disinfected with 70% ethyl alcohol. Then, skin scrapings and hair plucks were taken from the active margins of the lesions using sterile disposable scalpel blades as described previously [22]. The samples were transported to Central Veterinary Research Laboratory (CVRL)

located at Soba using clean, dry sterile Petri dish. Wet mounts were prepared from specimens with 20% potassium hydroxide for direct microscopic examination. Cultures were made onto duplicate sets of Sabouraud's dextrose agar (SDA) plates supplemented with 0.05 mg/ml chloramphenicol and 0.5 mg/ml cycloheximide and incubated aerobically at 27 and 37°C. Cultures were observed daily for growth of dermatophytes for up to four weeks. For more studies on cultural features, potato dextrose agar (PDA), cornmeal agar (Oxoid) and 5% horse blood agar were used. The plates showing no growth were discarded after 28 days of incubation while those showed growth were identified on the basis of their colonial and morphological characteristics including the growth rate, colony morphology; colour, shape, size and observe and reverse sides [8].

2.2.2 Biochemical study

Urease hydrolysis was done by inoculated parts of colonies from SDA on Christensen urea agar (Difco) to further confirm the identity of the isolated dermatophyte species [23].

2.2.3 Molecular study

Further, the isolated fungi were characterized by molecular markers using (β tubulin gene).

2.3.3.1 Molecular identification of isolates

2.3.3.1.1 DNA extraction and purification

The genomic DNA was extracted and purified from skin scrapings 10 days- old culture media following Möller *et al.*, method [24]. Briefly, mycelia were harvested from the surface of Sabouraud's dextrose agar culture, transferred to a mortar, supplemented with liquid nitrogen and ground to powder with a pestle. About 45 mg of the mycelia powder, was transferred to 2.0 ml micro tube, supplemented with 500 μ l TES (100 mM Tris, pH 8.0, 10 mM EDTA, 2% SDS) followed by addition of 3.75 μ l Proteinase K (stock 20mg/ml; w/v; Promega), before incubation for 1 hour at 60°C with occasional gentle shaking of the tube. The reaction mix was then supplemented with 140 μ l 5M NaCl, 700 μ l chloroform: isoamyl alcohol (24:1, v/v;) and incubated for 10 min at 65°C. Tubes were gently shaken, incubated for 30 min at 0°C and then centrifuged for 10 min in 4°C centrifuge

adjusted at 12,000 rpm. The supernatant was then transferred to 1.5 ml micro-tube, supplemented with 225 μ l 5 M NH_4Ac (Ammonium Acetate), mixed by gently finger mixing, placed on ice for one hour, before centrifugation at the same former conditions. The supernatant was transferred to a fresh 1.5 ml micro-tube, supplemented with 500 μ l isopropanol, placed on ice for 30 minutes, centrifuged for 5 min at the same former conditions. Finally the supernatant was decanted, the pellet washed twice with cold 70% ethanol, dried at room temperature, dissolved with 50 μ l TE (10 mM Tris-HCl, 1 mM disodium EDTA, pH 8.0) and stored at -20°C for later experiments.

2.3.3.1.2 Amplification of the β -tubulin gene PCR

Primers

A pair of primers, β t2a and β t2b synthesized by Invitrogen, were used [25]. Their sequences are

5'-GGTAACCAAATCGGTGCTGCTTTC-3' and 5'-ACCCTCAGTGTAGTGACCCTTGGC-3' respectively.

Reaction components

DreamTaq Green PCR Master Mix (2X) (Fermentas, Cat. # K1072) that contains all components in a volume of 25 μ l, except template DNA and primers was used. A number of four master mix tubes were used to make the reaction mix in a volume of 50 μ l; each tube received 1.5 μ l of each primer, 5 μ l of the respective template DNA and 17 μ l nuclease free H_2O .

Thermocycling conditions

The PCR was performed with a TC-512 (Techne) thermocycler following Samson, *et al.*, method [26]. The heat lid temperature was adjusted to 104°C. The thermocycling conditions were one cycle of 94°C for 1 min; 32 cycles of 94°C for 1 min, 68°C for 1 min and 72°C for 1 min; and one cycle of 72°C for 5 min.

Agarose gel

1.5% agarose (Vivantis) in 0.5x TBE (Tris Borate EDTA) supplemented with ethidium bromide (10 mg/ml, Vivantis) to a final concentration of 0.5 μ g/ml was used.

Electrophoresis

A Horizontal mini-gel electrophoresis system was used. One well was loaded with 5µl of 100 bp DNA size marker (Vivantis), and the other wells loaded with 8 µl of each PCR amplicon mixed with 2 µl of loading dye 6x (Vivantis). The power supply was adjusted to 80 V and the DNA was left to be separated for 45 min.

Visualization of resolved DNA

The separated DNA bands were visualized and photographed by a gel documentation system (Ingenius, Syngene Bio Imaging).

2.3.3.1.3 Sequencing and sequence analysis

The PCR amplicons were sent to JBMI (JEONJU BIOMATERIALS INSTITUTE) for automated bidirectional sequencing using the β t2a and β t2b primers. The sequences were analyzed by the BioEdit software package and the Basic Local Alignment Search Tool (BLAST) (<http://blast.ncbi.nlm.nih.gov>).

3. RESULTS

3.1 Case History

Infected animals showed skin lesions started as alopecia, severe incrustation, scaling, on the flanks backs, face, ears legs and gluteal areas similar to dermatophyte lesion. Other lesions of ringworm were observed on the withers and saddle area, but the infection was spread to the neck, chest and head with one or more legs being involved (Figs. 1 and 2).



Fig. 1. Localized skin lesions of dermatophytes on the face, neck and fore limbs

3.2 Morphological Characterization

3.2.1 Macroscopic characters

Two different isolated colonies were obtained on SDA, macroscopic feature of the isolated fungi for isolate 1 showed moderate rapid growth of a white to creamy flat powdery colony (Fig. 3). At 7 days on incubation, nodular granular white to creamy colony on obverse and a pale yellowish on reverse was observed (Fig. 4).



Fig. 2. Generalized skin lesions of dermatophytes



Fig. 3. White to creamy powdery colony of *T. mentagrophytes* on SDA

On potato dextrose agar and corn meal agar creamy colonies with granular surface on obverse and pale grey on reverse were obtained. On blood agar, as from days 5 a clear zone of β -haemolysis was observed around the colony.

Macroscopic feature for isolate 2 showed creamy, glabrous, flat, convoluted, lightly downy, heaped up with grayish reverse side, suggestive of *T. verrucosum*.



Fig. 4. Yellowish reverse colony of *T. mentagrophytes* on SDA




Fig. 5. Gel image of dermatophyte DNA amplified with β tubulin primers with a product size of approximately 500 bp

3.2.2 Microscopical investigation

On direct microscopic examination of hair and skin scrapings 19 samples showed fungal elements in the form of ectothrix arrangement of spores for some samples (isolate 1) and large numbers of endothrix spores and hyaline hyphae inside the hair for others (isolate 2). The microscopic feature (for isolate 1) showed numerous round micro conidia clustered on branched conidiophores and coiled hyphae. For isolate 2: Chains of chlamydo spores with antler hyphae were observed. Moreover, a positive urease reaction was developed on Christensen agar urea within 9 days for isolate 1 (*T. mentagrophytes*) and a negative one for isolate 2 (*T. verrucosum*).

3.3 Molecular Characterization

Analysis of the PCR amplicons by agarose gel electrophoresis revealed amplicon fragments of about 500bp as shown in Fig. 5.

3.3.1 Sequence analysis

3.3.1.1 Sequence analysis for *T. mentagrophytes* (isolate 1)

Analysis of the sequences of the isolate 1 using BioEdit software package showed that the clean sequence is 436 bp (Fig. 6). The sequence is unique and has similarity to the partial sequence of the beta-tubulin gene of many fungi including *T. mentagrophytes*. But the isolate was thoroughly investigated using the conventional mycological and biochemical assays that shown its similarity to *T. mentagrophytes*.

3.3.1.2 Sequence analysis for *T. verrucosum* (isolate 2)

Only 255 bp is clean sequence (Fig. 7). The clean sequence is similar to the partial sequence of the beta-tubulin gene of the *T. verrucosum* accession number KJ606180.1.

4. DISCUSSION

Donkeys (*Equus asinus*) are considered as important animals. They play an essential role in the economy of underdeveloped countries including Sudan [11]. In the present study, the clinical signs of alopecia, scaling and crusting, which appeared to varying degrees, suggested dermatophyte infection [8,27]. This finding was due to proteolytic and lipolytic enzymes that secreted by dermatophytes which favor digestion of skin tissue and hair resulting in hair loss, scaly and crusty lesions [5,8]. Furthermore, infection among equines showed that equine ringworm is highly contagious which transmitted by direct contact or indirect route through contaminated fomites. This indicates, possibility of existence of dermatophyte spores in the environment.

Mycological investigation revealed isolation of *T. mentagrophytes* and *T. verrucosum*. This finding is similar to that obtained [16,17,18] when investigated equines for skin infection.

Molecular characterization of the isolates was done on the basis that encouraged the use of PCR worldwide due to insensitivity of the conventional methods to detect fungal elements by direct microscopy and a non specificity of the

```

      10      20      30      40      50      60
GGTAACCAAA TCGGTGCTGC TTTCGCCTTC CCAGGCTTTG CTAATTGTGT CCACGTACAG 60
GCAAAACATC GCTGGTGAAC ATGGTCTCGA TGGTCCGGC ATGTAAGTGT GCTGGTTGAT 120
GGACATCTCC AACGATTCAC CTTTGACAAC CTTGAAACAG GTATGCTGGG TCTTCTGAAC 180
TCCAGCTTGA GCGTATGAAC GTTTATTTCA ACGAGGTGAA TTGAATTGG CAATATCCTA 240
CACTCAGTCC TTGCTTACCA CTCTCCAGGC CTCTGGCGGA AAATATGTCC CCCGTGCCGT 300
      310      320      330      340      350      360
CCTCGTCGAC CTTGAACCCG GTACCATGGA AGCCGTCAA GCTGGACCAT TCGGTGAACT 360
ATTCCGTTCC GATAACTTCG TCTTCGGACA ATCCGGTGCT GAAACAAC T GGGCCAAGG 420
TCACTACACT GAGGGT 436
    
```

Fig. 6. Partial sequence of *T. mentagrophytes* (isolate 1)

```

      10      20      30      40      50      60
TTGCAGCATA ATTGTATATC TCGTGTCAAT TGTACTGAC TTGATTGCA GGCAAACCAT 60
TGCCGGTGAG CACGGTCTCG ATGGATCCGG CCAGTGAGTG ATTCTGCAGG GGAGACAAAG 120
TCCCGGAGTC TCGAGGGACT TGAATGTGA CGATTGGGAT TTCTTTAGCT ACACCGGATC 180
TTCTGACCTC CAATTGGAGC GCATGAATGT CTACTTCAAC GAGGTGTGCA CGACCAAGAC 240
CCTTCCCTTC AGCAG 255
    
```

Fig. 7. Partial sequence of *T. verrucosum* (isolate 2)

culture. The conventional procedure is time-consuming usually requiring 10-15 days or even 3-4 weeks. Thus the use of molecular methods has supplemented traditional methods with DNA – based tools to examine phylogenetics and systematic of fungi [28]. This finding is similar to previous findings obtained [29] where re-classification of *Penicillium* species on molecular basis was performed. In the current study, the PCR technique used was found simple, inexpensive and rapid for efficiently isolated fungi. In addition, DNA was found suitable for use as a template for PCR amplification and sequences assay. Similar result was reported that shown the importance of the amount of extracted DNA as a parameter determining sensitivity of the PCR [30]. Moreover, the mycelium used in this study was directly recovered from Petri-dish cultures; the quality and quantity of DNA obtained were found suitable for molecular assays; it does not require the use of expensive and specialized equipment or hazardous reagents. Similar studies were carried to confirm dermatophytes infection recovered from culture [31,32]. Thus, PCR- based technique is recommended to be applied to dermatophytes recovered from environmental sources to increase sensitivity and specificity and the cost of diagnosis [33,34].

In the current study, the use of β - tubulin primers a&b to amplify the obtained DNA was similarly used [26] to determine sequences for 180 strains representing all accepted species of *Penicillium*.

Previous studies were carried with different primers to characterize Dermatophytes targeting various genes. (panDerm1 5'-GAAGAAGATTGTCGTTTGCATCGTCTC-3' and panDerm2 5'-CTCGAGGTCAAAGCACGCCAGAG-3') targets the chitin synthase-encoding gene (chitin synthase 1 – chs1) served for detection of dermatophytes in general, whereas (Trubrum-for 5'-TCTTTGAACGCACATTGCGCC-3' and Trubrum-rev 5'-CGGTCCTGAGGGCGCTGAA-3') targets internal transcribed spacer gene 2 (its2) for the specific detection of *T. rubrum* were used [35,36].

The obtained results indicated that incorporation of PCR technique in routine laboratory processing of skin scrapings augmented detection of dermatophytes. This finding is in accordance with the findings of other workers [35,36]. Furthermore, evaluation of the importance of the use of commercial PCR, to increase the detection rate of dermatophytes compared to culture alone was reviewed [37,38].

5. CONCLUSION

The present investigation showed that the integration of different methods and techniques led to identification of useful molecular marker for characterization of dermatophyte species. Thus, the study has aided the development of molecular marker to make identification of dermatophytes species more valuable.

CONSENT

It is not applicable.

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the authors.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Moriarty B, Hay R, Morris-Jones R. The diagnosis and management of tinea. *BMJ*. 2012;34:e4380.
2. Weitzman I, McGlennis MR. The genus *Arthroderma* and its later synonym *Nannizzia*. *Mycotaxon*. 1986;25:505-18.
3. Calderone RA. Immunoregulation of dermatophytosis. *J Crit Rev Microbiol*. 1989;16:339-684.
4. Rippon JW. Medical mycology: The pathogenic fungi and the pathogenic actinomycetes. 3rd Ed. Philadelphia: WB Saunders; 1988.
5. Hay RJ. Dermatophytosis and other superficial mycoses. In Principles and Practice of Infectious Diseases, 4th ed. Edited by G. L. Mandel, J. E. Bennett & R. Dolin. New York: Churchill Livingstone. 1995;261-283.
6. Abanmi A, Bakheshwain S, El Khizzi N, Zouman AR, Hantirah S, Al Harthi F, et al. Characteristics of superficial fungal infections in the Riyadh region of Saudi Arabia. *Int J Dermatol*. 2008;47(3):229-235.
7. Clayton Y, Midgley G. Identification of agents of superficial mycoses. In Evans EGV, Richardson MD, editors. *Medical Mycology: A Practical Approach* Oxford: IRL Press; 1989.
8. Padhye AA, Weitzman I. The dermatophytes. In: *Microbiology & Microbial Infections. Medical Mycology, Laboratory Diagnosis*, 1998;4(Chapter 13): 229-232.
9. Weitzman I, Summerbell RC. The dermatophytes. *J Clin Microbiol Rev*. 1995;8(2):240-59.
10. Shimozawa K, Anzai T, Kamada M, Takatori K. Fungal and bacterial isolation from race horses with infectious dermatosis. *J Equine Sci*. 1997;8(4):89-93.
11. Angara TE, Ibrahim A, Ismail A. The use of donkeys for transport: The case of Khartoum State, Sudan. *WIT Transactions on Ecology and the Environment*. 2011a;150:651-60.
12. Angara TE, Ibrahim A, Ismail A. The role of donkeys in income generation and the impact of endoparasites on their performance. *U. of K. J Vet Med and Anim prod*. 2011;2(2):65-89.
13. Fadlelmula A, Agab H, Lehorgene JM, Abbas B, Abdalla AED. First isolation of *Trichophyton verrucosum* as the etiology of ringworm in the Sudanese camels (*Camelus dromedarius*). *J Rev Elev Med Vet Pays Trops*. 1994;47(2):184-87.
14. Surendran K, Bhat RM, Bloor R, Nandakishore B, Sukumar D. A clinical and mycological study of dermatophytic infections. *Indian J Dermatol. Mycolo Round*. 2014;59(3):262-67.
15. Nardoni S, Rocchigiani G, Papini RA, Veneziano V, Brajon G, Martini M, et al. Dermatophytosis in donkeys (*Equus asinus*) due to *Microsporum racemosum* an unusual geophilic agent. *J Med Myco. L*. 2016;12:8-10.
16. Wisal GA, Elham AS, AbdoElGabar MA. A report on *Trichophyton verrucosum* in Donkeys in the Sudan. *The Sudan J Vet Res*. 2005;20:73-5.
17. Ali FE, Abu-Samra MT, Ibrahim AM. *Trichophyton mentagrophytes* infection in the domestic donkey (*Equus asinus asinus*). *J Ann Trop Med Parasitol*. 1981;75:623-626.
18. Fadlelmula A, Um El- Alim A. Ringworm in a horse caused by *Trichophyton verrucosum*. *Bull Anim Hlth Prod Afric*. 1983;33:17-18.
19. Quinn PJ, Markey RK. concise review of veterinary microbiology. Blackwell Oxford, United Kingdom; 2003.
20. Bagy MM, Abdel Mallek AY. Saprophytic and keratogenic fungi associated with animal's hair from Riyadh, Saudi Arabia. *Zbl Mikrobiol*. 1991;146:305-10.
21. Shathele MS. Mycological Studies of *Trichophyton mentagrophytes var. mentagrophytes* causing Ringworm in Horses in Al Ahsa Province, Kingdom of Saudi Arabia. *J Med Med Sci*. 2014; 5(8):157-61.

22. Cheesbrough M. Medical Laboratory Manual for Tropical Countries. 1st ed. Butterworth-Heinemann, Great Britain; 1992.
23. Campbell CK, Johnson EM, Philpot CM, Warnok DW. Identification of pathogenic fungi. Public Health Laboratory Service, London, United Kingdom; 1996.
24. Möller EM, Bahnweg G, Sandermann H, Geiger HH. A simple and efficient protocol for isolation of high molecular weight DNA from filamentous fungi, fruit bodies, and infected plant tissues. Nucleic Acids Res. 1992;(20) 22: 6115-6116.
25. Glass NL, Donaldson GC. Development of primer sets designed for use with PCR to amplify PCR conserved genes from filamentous Ascomycetes. J Appl Environ Microbiol. 1995;61:1323-29.
26. Samson KA, Kuijpers AFA, Houbraken Jos AMP, Frisvad JC. Phylogenetic analysis of *Penicillium* subgenus *Penicillium* using partial β - tubulin sequences. Studies in Mycology. 2004;49: 175-200.
27. Ural K, Yağc B, Ocal N. Cellular enzyme values in hunter/jumper and dressage horses with dermatophytosis. Arq Bras Med Vet Zootec. 2009;61:1233-37.
28. Bruns TD, White TJ, Taylor JW. Fungal molecular systematics. J Annu Rev Ecol Syst. 1991;22:525-64.
29. Bowman BH, Taylor JW, Brownlee AG, Lee J, Lu SD, Boysen TJ. Re classification of *Penicilliumroqueforti* Group into three Species on the basis of molecular genetics and Biochemical profiles. J Microbiol. 1996;142:541-49.
30. Bontems O, Hauser PM, Monod M. Evaluation of a polymerase chain reaction-restriction fragment length polymorphism assay for dermatophyte and nondermatophyte identification in onychomycosis. Br J Dermatol. 2009;161: 791-96.
31. Jenssen RH, Arendrup MC. Molecular diagnosis of dermatophyte infection. J Curr Opin infec Dis. 2012; 25:126-34.
32. Anastasia S, Christina B, Eleni J, Evangelos DA, Myrto C. Evaluation of a commercial PCR test for the diagnosis of dermatophyte nail infections. J Med Microbiol. 2015;64:25-31.
33. Bergman A, Heimer D, Kondori N, Enroth H. Fast and specific dermatophyte detection by automated DNA extraction and real-time PCR. J Clin Microbiol Infect. 2013;19:205-11.
34. Brasch J, Beck-Jendroschek V, Glaser R. Fast and sensitive detection of *Trichophyton rubrum* superficial tinea and onychomycosis by use of a direct polymerase chain reaction assay. Mycoses. 2011;54:313–17.
35. Brillowska-Dabrowska A, Saunte DM, Arendrup MC. Five-hour diagnosis of dermatophyte nail infections with specific detection of *Trichophyton rubrum*. J Clin Microbiol. 2007;45:1200-04.
36. Brillowska-Dabrowska A, Nielsen SS, Nielsen HV, Arendrup MC. Optimized 5-hour multiplex PCR test for the detection of tinea unguium: performance in a routine PCR laboratory. J Med Mycol. 2010;48: 828–31.
37. Chandran NS, pan JY, Pramono ZA, Tan HH, Seow CS. Complementary role of a polymerase chain reaction test in the diagnosis of onychomycosis. Australas J Dermatol. 2013;54:105-108.
38. Kondori N, Abrahamsson AL, Ataollahy N, Wenneras C. Comparison of a new commercial test, Dermatophyte-PCR kit, with conventional methods for rapid detection and identification of *Trichophyton rubrum* in nail specimens. J Med Mycol. 2010;48:1005-08.

© 2017 Suleiman et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://sciedomain.org/review-history/21370>